

Date Mailed: March 22, 2004

Ser. No. 10/634,477 Filed: Aug. 4, 2003


Patent No. _____

Issued _____

Inventor(s) Paul Lehmann et al.

Title Treatment of disturbances of cross
distribution

- ☐ Assignment
- ☐ Change of Correspondence Address
- ☐ Fee Address Indication Form
- ☐ Issue Fee
- ☒ Priority Document
- ☐ Other: _____



Ref. No. 21368 US BRT: bah

BEST AVAILABLE COPY



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02019100.3

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

**CERTIFIED COPY OF
PRIORITY DOCUMENT**

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 02019100.3
Demande no:

Anmeldetag:
Date of filing: 29.08.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

F. HOFFMANN-LA ROCHE AG

4070 Basel
SUISSE *

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se referer à la description.)

Use of erythropoietin

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A61K38/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

Case 21368

USE OF ERYTHROPOIETIN

The present invention relates to a new use of erythropoietin, especially the treatment of disturbances of iron distribution in diabetes.

Various diseases are known, in which the metabolism of iron is not normal. In an
5 anemia, not enough blood can be formed due to an overall lack of iron in the body. An
other metabolic condition relating to iron is hemochromatosis, in which the overall
concentration of iron in the body is higher than normal, which leads to various conditions
such as e.g. the destruction of organs.

Disturbances of iron distribution differ from the above described anemia and
10 hemochromatosis because the overall concentration of iron in the body is normal. On one
hand, iron is accumulated in various organs and can lead to damages and even destruction
of these organs. On the other hand the use of the iron which is present in normal
quantities in the formation of blood is impaired, leading to secondary effects which are
comparable to those related to anemia.

15 Until now it was not known that patients suffering from diabetes have a high
probability to be affected by disturbances of iron distribution. Disturbances of iron
distribution can be diagnosed by various parameters which are commonly used in the
diagnosis of the iron status. Based on measurements of ferritin and soluble transferrin
receptor it is possible to assess whether the overall concentration of iron in a diabetic
20 patient is normal. If this is the case, then a lowered concentration of Hemoglobin in
reticulocytes is an indicator for disturbances of iron distribution. Another indicator is a
continuously/prolonged elevated concentration of C-reactive protein (CRP) in patients
suffering from diabetes and exhibiting a normal overall iron concentration. A method for

diagnosing disturbances of iron distribution has been described by P. Lehmann, M. Volkmann, J. Lotz, A. Baldauf, R. Roeddiger, poster presented at the AACC/CSCC, Annual Meeting, July 29 – August 2, 2001, Chicago, Illinois.

5 So far, no treatment has yet been suggested for diabetic patients suffering from disturbances in iron distribution. The problem underlying the present invention is therefore to provide a treatment for disturbances of iron distribution in diabetes in order to minimize or suppress the above mentioned disadvantages. It has surprisingly been found that erythropoietin has a beneficial effect on disturbances of iron distribution in diabetes. The problem is therefore solved, according to the present invention, by providing
10 erythropoietin for the use in the treatment of disturbances of iron distribution in diabetes.

Unless otherwise indicated the following definitions are set forth to illustrate and define the meaning and scope of the various terms used to describe the invention herein.

The term "lower-alkyl" as used herein means a linear or branched alkyl group having from one to six carbon atoms. Examples of lower-alkyl groups include methyl, ethyl and
15 isopropyl, preferably methyl.

The term "lower-alkoxy" as used herein means a group R'-O-, wherein R' is a lower-alkyl as described above.

The term "disturbances of iron distribution in diabetes" refers to a disturbance of iron distribution which occurs in patients suffering from diabetes. The disturbance of iron
20 distribution can e.g. be characterised as described above. Particularly, a disturbance of iron distribution is characterised by the following parameters: concentration of soluble transferrin receptor [mg/L] divided by log(concentration of ferritin [µg/L]) is smaller than 3.5 and simultaneously concentration of C-reactive protein is above 5 mg/L.

The term "erythropoietin" or "erythropoietin protein" refers to a protein with the *in*
25 *vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells and selected from the group consisting of human erythropoietin and analogs which are defined below.

The term "pegylated erythropoietin (Peg-EPO or PEG-EPO)" refers to an erythropoietin protein which is covalently linked with one to three polyethylene
30 derivatives as described below.

Description of Drawings:

Figure 1: Primary structure of human EPO (165 amino acids) (SEQ ID NO:1).

Figure 2: Primary structure of human EPO (166 amino acids) (SEQ ID NO:2).

In more detail, the present invention relates to the use of erythropoietin in the
5 manufacture of a medicament for the treatment of disturbances of iron distribution in
diabetes. In a preferred embodiment, the invention relates to a use as defined above,
wherein diabetes is non insulin dependant diabetes mellitus.

The present invention is especially useful for the preparation of pharmaceutical
compositions comprising erythropoietin as pharmaceutically active ingredient. The term
10 "erythropoietin" or "erythropoietin protein" or "EPO" is as follows: particularly the terms
refer to a glycoprotein, e.g. the human erythropoietin, e.g. having the amino acid sequence
set out in (SEQ ID NO: 1) or (SEQ ID NO: 2) or an amino acid sequence substantially
homologous thereto, whose biological properties relate to the stimulation of red blood cell
production and the stimulation of the division and differentiation of committed erythroid
15 progenitors in the bone marrow. As used herein, these terms include such proteins
modified deliberately, as for example, by site directed mutagenesis or accidentally through
mutations. These terms also include analogs having from 1 to 6 additional sites for
glycosylation, analogs having at least one additional amino acid at the carboxy terminal
end of the glycoprotein, wherein the additional amino acid includes at least one
20 glycosylation site, and analogs having an amino acid sequence which includes a
rearrangement of at least one site for glycosylation. These terms include both natural and
recombinantly produced human erythropoietin. In a preferred embodiment of the present
invention, the erythropoietin protein is a human erythropoietin.

As set out in detail below, the preparation and purification of EPO are well known in
25 the art. By erythropoietin is meant the natural or recombinant protein, preferably human,
e.g. epoetin alfa or epoetin beta, as obtained from any conventional source such as tissues,
protein synthesis, cell culture with natural or recombinant cells. Any protein having the
activity of erythropoietin, such as muteins or otherwise modified proteins, is
encompassed. In a preferred embodiment of the present invention, the erythropoietin
30 protein is epoetin alfa or epoetin beta. Recombinant EPO may be prepared via expression
in CHO-, BHK- or HeLa cell lines, by recombinant DNA technology or by endogenous
gene activation. Expression of proteins, including, by endogenous gene activation, is well

known in the art and is disclosed, for example in U.S. Patents Nos. 5,733,761, 5,641,670, and 5,733,746, and international patent publication Nos. WO 93/09222, WO 94/12650, WO 95/31560, WO 90/11354, WO 91/06667 and WO 91/09955, the contents of each of which are incorporated herein by reference. The use as defined above, wherein the

5 erythropoietin protein is expressed by endogenous gene activation, is preferred. The preferred EPO species for the preparation of erythropoietin glycoprotein products are human EPO species. More preferably, the EPO species is the human EPO having the amino acid sequence set out in SEQ ID NO:1 or SEQ ID NO:2, more preferably the amino acid sequence SEQ ID NO:1. A preferred embodiment of the present invention therefore

10 relates to the use as described above, wherein the erythropoietin protein has the amino acid sequence of SEQ ID NO:1 or SEQ ID NO:2.

Further, erythropoietin may be a glycoprotein analog having from 1 to 6 additional sites for glycosylation. Therefore, the present invention also relates to the use as described before, wherein the erythropoietin protein has the sequence of human erythropoietin

15 modified by the addition of from 1 to 6 glycosylation sites. Glycosylation of a protein, with one or more oligosaccharide groups, occurs at specific locations along a polypeptide backbone and greatly affects the physical properties of the protein such as protein stability, secretion, subcellular localization, and biological activity. Glycosylation is usually of two types. O-linked oligosaccharides are attached to serine or threonine residues and N-linked

20 oligosaccharides are attached to asparagine residues. One type of oligosaccharide found on both N-linked and O-linked oligosaccharides is N-acetylneuraminic acid (sialic acid), which is a family of amino sugars containing 9 or more carbon atoms. Sialic acid is usually the terminal residue on both N-linked and O-linked oligosaccharides and, because it bears a negative charge, confers acidic properties to the glycoprotein. Human erythropoietin,

25 having 165 amino acids, contains three N-linked and one O-linked oligosaccharide chains which comprise about 40% of the total molecular weight of the glycoprotein. N-linked glycosylation occurs at asparagine residues located at positions 24, 38, and 83 and O-linked glycosylation occurs at a serine residue located at position 126. The oligosaccharide chains are modified with terminal sialic acid residues. Enzymatic removal of all sialic acid

30 residues from the glycosylated erythropoietin results in loss of *in vivo* activity but not *in vitro* activity because sialylation of erythropoietin prevents its binding, and subsequent clearance, by hepatic binding protein.

The term "erythropoietin" includes analogs of human erythropoietin with one or more changes in the amino acid sequence of human erythropoietin which result in an

increase in the number of sites for sialic acid attachment. These glycoprotein analogs may be generated by site-directed mutagenesis having additions, deletions, or substitutions of amino acid residues that increase or alter sites that are available for glycosylation. Glycoprotein analogs having levels of sialic acid greater than those found in human erythropoietin are generated by adding glycosylation sites which do not perturb the secondary or tertiary conformation required for biological activity. The glycoproteins of the present invention also include analogs having increased levels of carbohydrate attachment at a glycosylation site which usually involve the substitution of one or more amino acids in close proximity to an N-linked or O-linked site. The glycoproteins of the present invention also include analogs having one or more amino acids extending from the carboxy terminal end of erythropoietin and providing at least one additional carbohydrate site. The erythropoietin proteins of the present composition also include analogs having an amino acid sequence which includes a rearrangement of at least one site for glycosylation. Such a rearrangement of glycosylation site involves the deletion of one or more glycosylation sites in human erythropoietin and the addition of one or more non-naturally occurring glycosylation sites. Increasing the number of carbohydrate chains on erythropoietin, and therefore the number of sialic acids per erythropoietin molecules may confer advantageous properties such as increased solubility, greater resistance to proteolysis, reduced immunogenicity, increased serum half-life, and increased biological activity. Erythropoietin analogs with additional glycosylation sites are disclosed in more detail in European Patent Application 640 619, to Elliot published March 1, 1995.

In a preferred embodiment, the pharmaceutical composition of the present invention comprises erythropoietin proteins with an amino acid sequence which includes at least one additional site for glycosylation such as, but not limited to, erythropoietins comprising the sequence of human erythropoietin modified by a modification selected from the following:

Asn³⁰Thr³²;
Asn⁵¹Thr⁵³;
Asn⁵⁷Thr⁵⁹;
Asn⁶⁹;
Asn⁶⁹Thr⁷¹;
Ser⁶⁸Asn⁶⁹Thr⁷¹;
Val⁸⁷Asn⁸⁸Thr⁹⁰;
Ser⁸⁷Asn⁸⁸Thr⁹⁰;
Ser⁸⁷Asn⁸⁸Gly⁸⁹Thr⁹⁰;

Ser⁸⁷ Asn⁸⁸ Thr⁹⁰ Thr⁹²;
 Ser⁸⁷ Asn⁸⁸ Thr⁹⁰ Ala¹⁶²;
 Asn⁶⁹ Thr⁷¹ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰;
 Asn³⁰ Thr³² Val⁸⁷ Asn⁸⁸ Thr⁹⁰;
 5 Asn⁸⁹ Ile⁹⁰ Thr⁹¹;
 Ser⁸⁷ Asn⁸⁹ Ile⁹⁰ Thr⁹¹;
 Asn¹³⁶ Thr¹³⁸;
 Asn¹³⁸ Thr¹⁴⁰;
 Thr¹²⁵; and
 10 Pro¹²⁴ Thr¹²⁵.

The notation used herein for modification of amino acid sequence means that the position(s) of the corresponding unmodified protein (e.g. hEPO of SEQ ID NO:1 or SEQ ID NO:2) indicated by the superscripted number(s) is changed to the amino acid(s) that immediately precede the respective superscripted number(s).

15 The erythropoietin protein may also be an analog having at least one additional amino acid at the carboxy terminal end of the glycoprotein, wherein the additional amino acid includes at least one glycosylation site. The additional amino acid may comprise a peptide fragment derived from the carboxy terminal end of human chorionic gonadotropin. Preferably, the glycoprotein is an analog selected from the group consisting
 20 of (a) human erythropoietin having the amino acid sequence, Ser Ser Ser Ser Lys Ala Pro Pro Pro Ser Leu Pro Ser Pro Ser Arg Leu Pro Gly Pro Ser Asp Thr Pro Ile Leu Pro Gln, extending from the carboxy terminus; (b) the analog in (a) further comprising Ser⁸⁷ Asn⁸⁸ Thr⁹⁰ EPO; and (c) the analog in (a) further comprising Asn³⁰ Thr³² Val⁸⁷ Asn⁸⁸ Thr⁹⁰ EPO.

25 The erythropoietin protein may also be an analog having an amino acid sequence which includes a rearrangement of at least one site for glycosylation. The rearrangement may comprise a deletion of any of the N-linked carbohydrate sites in human erythropoietin and an addition of an N-linked carbohydrate site at position 88 of the amino acid sequence of human erythropoietin. Preferably, the glycoprotein is an analog
 30 selected from the group consisting of Gln²⁴ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰ EPO; Gln³⁸ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰ EPO; and Gln⁸³ Ser⁸⁷ Asn⁸⁸ Thr⁹⁰ EPO. A further analog is darbepoetin alfa. A preferred erythropoietin protein in the use described before is darbepoetin alfa.

More particularly, the erythropoietin protein of the present pharmaceutical composition as described above may also include pegylated derivatives thereof. Pegylated derivatives of erythropoietin and their preparation are known in the art and described for example in WO 01/02017, EP-A-1064951, EP-A-539,167, EP-A-605,963, WO 93/25212, WO 94/20069, WO 95/11924, US Patent No. 5,56, EP-A-584,876, WO 92/16555, WO 94/28024, WO 97/04796, US Pat. Nos. 5,359,030 and 5,681,811, US Patent No. 4,179,337, Japanese Patent, WO 98/32466, US Patent No. 5,324,650. Preferable, in the use described above, the erythropoietin protein is pegylated. A preferred embodiment of pegylated erythropoietin species refer to the derivatives as described below.

Accordingly, the present invention also refers to the use as described above, wherein the erythropoietin protein is a conjugate, said conjugate comprising an erythropoietin protein as described above having at least one free amino group and having the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells and selected from the group consisting of human erythropoietin and analogs thereof which have a sequence of human erythropoietin modified by the addition of from 1 to 6 glycosylation sites or a rearrangement of at least one glycosylation site; said erythropoietin being covalently linked to n poly(ethylene glycol) groups of the formula $\text{CO}-(\text{CH}_2)_x-(\text{OCH}_2\text{CH}_2)_m-\text{OR}$ with the CO (i.e. carbonyl) of each poly(ethylene glycol) group forming an amide bond with one of said amino groups; wherein R is lower-alkyl; x is 2 or 3; m is from about 450 to about 900; n is from 1 to 3; and n and m are chosen so that the molecular weight of the conjugate minus the erythropoietin protein is from 20 kilodaltons to 100 kilodaltons. This invention further provides pharmaceutical compositions containing conjugates described herein in which the percentage of conjugates in which n is 1 is at least ninety percent, preferably at least ninety-two percent, ore preferably ninety-six percent of all conjugates of the composition.

More specifically the above conjugates may be represented by formula (I)




wherein P is the residue of an erythropoietin protein as described herein, (i.e. without the amino group or amino groups which form an amide linkage with the carbonyl shown in Formula I), having the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells; and wherein R is lower-alkyl; x is 2 or 3; m is from about 450 to about 900; n is from 1 to 3; and n and m are chosen so that the molecular weight of the conjugate minus the erythropoietin glycoprotein is from 20

kilodaltons to 100 kilodaltons. In accordance with this invention, R is any lower-alkyl. Conjugates in which R is methyl are preferred.

The symbol "m" represents the number of ethylene oxide residues (OCH₂CH₂) in the poly(ethylene oxide) group. A single PEG (polyethylene glycol) subunit of ethylene
 5 oxide has a molecular weight of about 44 daltons. Thus, the molecular weight of the conjugate (excluding the molecular weight of the EPO) depends on the number "m". In the conjugates of this invention "m" is from about 450 to about 900 (corresponding to a molecular weight of about 20 kDa to about 40 kDa), preferably from about 650 to about 750 (corresponding to a molecular weight of about 30 kDa). The number m is selected
 10 such that the resulting conjugate of this invention has a physiological activity comparable to unmodified EPO, which activity may represent the same as, more than, or a fraction of the corresponding activity of unmodified EPO. A molecular weight of "about" a certain number means that it is within a reasonable range of that number as determined by conventional analytical techniques. The number "m" is selected so that the molecular
 15 weight of each poly(ethylene glycol) group covalently linked to the erythropoietin glycoprotein is from about 20kDa to about 40kDa, and is preferably about 30 kDa.

In the conjugates of this invention, the number "n is the number of poly(ethylene glycol) groups covalently bound to free amino groups (including ε-amino groups of a lysine amino acid and/or the amino-terminal amino group) of an erythropoietin protein
 20 via amide linkage(s). A conjugate of this invention may have one, two, or three PEG groups per molecule of EPO. "n" is an integer ranging from 1 to 3, preferably "n" is 1 or 2, and more preferably "n" is 1. A preferred conjugate of the conjugates described above comprises compounds wherein x is 2, m is 650 to 750, n is 1 and R is methyl.

The compound of formula (I) can be prepared from the known polymeric material:




25

in which R and m are as described above, by condensing the compound of Formula II with the erythropoietin glycoprotein. Compounds of formula (II) in which x is 3 are alpha-lower-alkoxy, butyric acid succinimidyl esters of poly(ethylene glycol) (lower-alkoxy-PEG-SBA). Compounds of formula (II) in which x is 2 are alpha-lower-alkoxy, propionic acid

succinimidyl esters of poly(ethylene glycol) (lower-alkoxy-PEG-SPA). Any conventional method of reacting an activated ester with an amine to form an amide can be utilized. In the reaction described above, the exemplified succinimidyl ester is a leaving group causing the amide formation. The use of succinimidyl esters such as the compounds of formula II
5 to produce conjugates with proteins are disclosed in U.S. Patent No. 5,672,662, issued September 30, 1997 (Harris, et al.).


Human EPO contains nine free amino groups, the amino-terminal amino group plus the ϵ -amino groups of 8 lysine residues. When the pegylation reagent was combined with a SBA compound of Formula II, it has been found that at pH 7.5, a protein:PEG ratio
10 of 1:3, and a reaction temperature of from 20-25 °C, a mixture of mono-, di-, and trace amounts of the tri-pegylated species were produced. When the pegylation reagent was a SPA compound of Formula II, at similar conditions except that the protein:PEG ratio was 1:2, primarily the mono-pegylated species is produced. The pegylated EPO can be administered as a mixture, or as the cation exchange chromatography separated different
15 pegylated species. By manipulating the reaction conditions (e.g., ratio of reagents, pH, temperature, protein concentration, time of reaction etc.), the relative amounts of the different pegylated species can be varied.

A further preferred embodiment of the present invention relates to the use as defined above, wherein the erythropoietin protein is a conjugate, said conjugate comprising an
20 erythropoietin protein as defined above having at least one free amino group and having the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells and selected from the group consisting of human erythropoietin protein and analogs thereof which have the primary structure of human erythropoietin protein modified by the addition of from 1 to 6 glycosylation sites; said
25 erythropoietin protein being covalently linked to from one to three lower-alkoxy poly(ethylene glycol) groups, each poly(ethylene glycol) group being covalently linked to the erythropoietin protein *via* a linker of the formula $-C(O)-X-S-Y-$ with the C(O) of the linker forming an amide bond with one of said amino groups, X is $-(CH_2)_k-$ or $-CH_2(O-CH_2-CH_2)_k-$, k is from 1 to 10, Y is




the average molecular weight of each poly(ethylene glycol) moiety is from about 20 kilodaltons to about 40 kilodaltons, and the molecular weight of the conjugate is from about 51 kilodaltons to about 175 kilodaltons.

5 This erythropoietin species may also be represented by formula (III)




wherein R may be any lower-alkyl. A preferred lower-alkyl is methyl. X may be $-(CH_2)_k-$ or $-CH_2(O-CH_2-CH_2)_k-$, wherein k is from 1 to about 10. Preferably, k is from 1 to about 4, more preferably, k is 1 or 2. Most preferably, X is $-(CH_2)$.


10 In Formula 1, Y is



preferably




more preferably




In formula (III), the number m is selected such that the resulting conjugate of formula (III) has a physiological activity comparable to unmodified EPO, which activity may represent the same as, more than, or a fraction of the corresponding activity of unmodified EPO. m represents the number of ethylene oxide residues in the PEG unit. A single PEG subunit of $-(OCH_2CH_2)-$ has a molecular weight of about 44 daltons. Thus, the molecular weight of the conjugate (excluding the molecular weight of the EPO) depends on the number m . A molecular weight of "about" a certain number means that it is within a reasonable range of that number as determined by conventional analytical techniques. m is an integer ranging from about 450 to about 900 (corresponding to a molecular weight of from 20 to 40 kDa), preferably m is from about 550 to about 800 (about 24 to 35 kDa), and most preferably m is from about 650 to about 700 (about 29 to about 31 kDa).

In formula (III), the number n is the number of ϵ -amino groups of a lysine amino acid in an erythropoietin protein covalently bound to a PEG unit via an amide linkage. A conjugate of this invention may have one, two, or three PEG units per molecule of EPO. n is an integer ranging from 1 to 3, preferably n is 1 or 2, and more preferably n is 1.


Preferred erythropoietin proteins of formula (III) are represented by the formulae:



and





In the most preferred embodiment of the present invention, an erythropoietin conjugate is represented by the formula:




wherein in the above formulae n is an integer from 1 to 3; m is an integer from 450 to 900; R is lower-alkyl; X is $-(\text{CH}_2)_k-$ or $-\text{CH}_2(\text{O}-\text{CH}_2-\text{CH}_2)_k-$, and P is the residue of the erythropoietin protein without the amino group or groups which form an amide linkage
5 with X .

Other preferred erythropoietin glycoprotein products are represented by the formulae:



More preferred erythropoietin glycoprotein products are represented by the formula:



10


These erythropoietin proteins may be prepared by

(a) covalently reacting an ϵ -amino group of a lysine amino acid of an erythropoietin protein represented by the formula, $\text{P}-[\text{NH}_2]_n$, with a bi-functional reagent represented by the formula, $\text{Z}-\text{CO}-\text{X}-\text{S}-\text{Q}$, to form an intermediate with an amide linkage represented by
15 the formula:



wherein P is an erythropoietin protein less the amino group which forms an amide linkage; n is an integer ranging from 1 to 3; Z is a reactive group, e.g. a carboxylic-NHS ester; X is $-(CH_2)_k-$ or $-CH_2(O-CH_2-CH_2)_k-$, wherein k is from 1 to about 10; and Q is a protecting group, like alkanoyl, e.g. acetyl.


- 5 (b) covalently reacting the intermediate with an amide linkage from step (a) with an activated poly(ethylene glycol) derivative represented by the formula, $W-[OCH_2CH_2]_m-OR$, to form an erythropoietin glycoprotein product represented by the formula:



- 10 wherein W is a sulfhydryl reactive form of Y; m is an integer ranging from about 450 to about 900; R is lower-alkyl; and Y is as defined above.


- In this embodiment, the bi-functional reagent is preferably N-succinimidyl-S-acetylthiopropionate or N-succinimidyl-S-acetylthioacetate, Z is preferably N-hydroxy-succinimide, and the activated poly(ethylene glycol) derivative $W-[OCH_2CH_2]_m-OR$ is preferably selected from the group consisting of iodo-acetyl-methoxy-PEG, methoxy-PEG-15 vinylsulfone, and methoxy-PEG-maleimide.

- In more detail, the erythropoietin proteins of formula (III) may be prepared by covalent linking of thiol groups to EPO ("activation") and coupling the resulting activated EPO with a poly(ethylene glycol) (PEG) derivative. The first step for the preparation of pegylated EPO according to the present invention comprises covalent linking of thiol
20 groups via NH_2 -groups of EPO. This activation of EPO is performed with bi-functional reagents which carry a protected thiol group and an additional reactive group, such as active esters (e.g., a succinimidylester), anhydrides, esters of sulphonic acids, halogenides of carboxylic acids and sulphonic acids, respectively. The thiol group is protected by groups known in the art, e.g., acetyl groups. These bi-functional reagents are able to react
25 with the ξ -amino groups of the lysine amino acids by forming an amide linkage. The first step of the reaction is set out below:




EPO, n and X are as defined above and Z is a reactive group known in the art, e.g. a N-hydroxy-succinimide (NHS) substituent of the formula


5




In a preferred embodiment the activation of the ϵ -amino lysine groups is performed by reaction with bi-functional reagents having a succinimidyl moiety. The bi-functional reagents may carry different spacer species, e.g. $-(\text{CH}_2)_k-$ or $-\text{CH}_2-(\text{O}-\text{CH}_2-\text{CH}_2-)_k-$ moieties, wherein k is from 1 to about 10, preferably from 1 to about 4, and more preferably 1 or 2, and most preferably 1. Examples of these reagents are N-succinimidyl-S-acetylthiopropionate (SATP) and N-succinimidyl-S-acetylthioacetate (SATA)




Acetylthioalkyl-carboxylic-NHS-ester, like



SATP



SATA



2-(Acetylthio)-(ethoxy)_k-acetic-acid-NHS-ester

with k as defined above.


The preparation of the bi-functional reagents is known in the art. Precursors of 2-(acetylthio)-(ethoxy)_k-acetic-acid-NHS-esters are described in DE-3924705, while the derivatization to the acetylthio compound is described by March, J., Advanced Organic Chemistry, McGraw-Hill, 1977, 375-376. SATA is commercially available (Molecular Probes, Eugene, OR, USA and Pierce, Rockford, IL).

The number of thiol groups to be added to an EPO molecule can be selected by adjusting the reaction parameters, i.e., the protein (EPO) concentration and the protein/bi-functional reagent ratio. Preferably, the EPO is activated by covalently linking from 1 to 5 thiol groups per EPO molecule, more preferably from 1.5 to 3 thiol groups per EPO molecule. These ranges refer to the statistical distribution of the thiol group over the EPO protein population.

The reaction is carried out, for example, in an aqueous buffer solution, pH 6.5–8.0, e.g., in 10 mM potassium phosphate, 50 mM NaCl, pH 7.3. The bi-functional reagent may be added in DMSO. After completion of the reaction, preferably after 30 minutes, the reaction is stopped by addition of lysine. Excess bifunctional reagent may be separated by methods known in the art, e.g., by dialysis or column filtration. The average number of thiol groups added to EPO can be determined by photometric methods described in, for example, Grasetti, D.R. and Murray, J.F. in J. Appl. Biochem. Biotechnol. 119, 41 – 49 (1967).


The above reaction is followed by covalent coupling of an activated poly(ethylene glycol) (PEG) derivative. Suitable PEG derivatives are activated PEG molecules with an average molecular weight of from about 20 to about 40 kDa, more preferably from about 24 to about 35 kDa, and most preferably about 30 kDa.

Activated PEG derivatives are known in the art and are described in, for example, Morpurgo, M. et al. J. Bioconj. Chem. (1996) 7, page 363 ff for PEG-vinylsulfone. Linear chain and branched chain PEG species are suitable for the preparation of the compounds of Formula 1. Examples of reactive PEG reagents are iodo-acetyl-methoxy-PEG and methoxy-PEG-vinylsulfone:




The use of these iodo-activated substances is known in the art and described e.g. by Hermanson, G. T. in *Bioconjugate Techniques*, Academic Press, San Diego (1996) p. 147-148.

Most preferably, the PEG species are activated by maleimide using (lower-alkoxy-
5 PEG-maleimide), such as methoxy-PEG-maleimide (MW 30000; Shearwater Polymers, Inc.). The structure of lower-alkoxy-PEG-maleimide is as follows:



with R and m are as defined above, preferably



10 The coupling reaction with lower-alkoxy-PEG-maleimide takes place after *in situ* cleavage of the thiol protecting group in an aqueous buffer solution, e.g. 10 mM potassium phosphate, 50 mM NaCl, 2 mM EDTA, pH 6.2. The cleavage of the protecting group may be performed, for example, with hydroxylamine in DMSO at 25°C, pH 6.2 for about 90
15 minutes. For the PEG modification the molar ratio of activated EPO/lower-alkoxy-PEG-maleimide should be from about 1:3 to about 1:6, and preferably 1:4. The reaction may be stopped by addition of cysteine and reaction of the remaining thiol (-SH) groups with N-methylmaleimide or other appropriate compounds capable of forming disulfide bonds. Because of the reaction of any remaining active thiol groups with a protecting group such as N-methylmaleimide or other suitable protecting group, the EPO glycoproteins in the
20 conjugates of this invention may contain such protecting groups. Generally the procedure described herein will produce a mixture of molecules having varying numbers of thiols protected by different numbers of the protecting group, depending on the number of activated thiol groups on the glycoprotein that were not conjugated to PEG-maleimide.

Whereas N-methylmaleimide forms the same type of covalent bond when used to block the remaining thiol-groups on the pegylated protein, disulfide compounds will lead in an intermolecular sulfide/disulfide exchange reaction to a disulfide bridged coupling of the blocking reagent. Preferred blocking reagents for that type of blocking reaction are oxidized glutathione (GSSG), cysteine and cystamine. Whereas with cysteine no additional net charge is introduced into the pegylated protein, the use of the blocking reagents GSSG or cystamine results in an additional negative or positive charge.

The further purification of the compounds of formula (III), including the separation of mono-, di- and tri-pegylated EPO species, may be done by methods known in the art, e.g., column chromatography.

Pegylated erythropoietin derivatives preferably contains at least ninety percent mono-PEG conjugates. Usually mono-PEG conjugates of erythropoietin glycoproteins are desirable because they tend to have higher activity than di-PEG conjugates.. The percentage of mono-PEG conjugates as well as the ratio of mono- and di-PEG species can be controlled by pooling broader fractions around the elution peak to decrease the percentage of mono-PEG or narrower fractions to increase the percentage of mono-PEG in the composition. About ninety percent mono-PEG conjugates is a good balance of yield and activity. Sometimes compositions in which, for example, at least ninety-two percent or at least ninety-six percent of the conjugates are mono-PEG species (n equals 1) may be desired. In an embodiment of this invention the percentage of conjugates where n is 1 is from ninety percent to ninety-six percent.

Pharmaceutical compositions comprising pegylated erythropoietin are known in the art and are described e.g. in International patent application WO 01/87329. Compositions may comprise 10 to 10000 µg of an erythropoietin protein per ml as defined above. Preferably, the compositions comprise 10 to 1000 µg, e.g. 10, 50, 100, 400, 800 or 2500 µg per ml. Further, the compositions may comprise 10 µg to 10000 µg erythropoietin protein per ml, 10 – 200 mmol/l sulfate, 10 to 50 mmol/l phosphate, pH 6.0 to 6.5. This composition may also comprise up to 20 mM methionine, 1 – 5 % of a polyol (w/v), up to 0.1 % pluronic F68 (w/v) and optionally up to 1 mM CaCl₂. An example of this composition comprises 10 µg to 10000 µg erythropoietin protein per ml, 40 mmol/l sulfate, 10 mmol/l phosphate, 3% mannitol (w/v), 10 mM methionine, 0.01% pluronic F68 (w/v), pH 6.2. In alternative the composition may comprise 10 µg to 10000 µg erythropoietin protein per ml, 10 to 100 mmol/l NaCl, 10 to 50 mmol/l phosphate pH 6.0 to 7.0, optionally 1-5% (w/v) of a polyol. Further, this composition may comprise up to 20

mM methionine, up to 0.1% pluronic F68 (w/v) and optionally 7.5 $\mu\text{mol/l}$ CaCl_2 . Specifically, this composition may comprise 10 μg to 10000 μg erythropoietin protein per ml, 100 mmol/l NaCl, 10 mM methionine, 0.01% pluronic F68 (w/v), and 10 mmol/l phosphate, pH 7.0.

5 The present invention also refers to the above composition comprising 10 μg to 10000 μg erythropoietin protein per ml, 10 to 50 mmol/l arginine, pH 6 to pH 6.5, 10 to 100 mmol/l sodium sulfate. In addition, this composition may comprise up to 20 mM methionine, up to 0.1% pluronic F68 (w/v), optionally up to 1 mmol/l CaCl_2 and
° optionally 1 – 5 % (w/v) of a polyol. Specifically, this composition may 10 μg to 10000 μg
10 erythropoietin protein per ml, 40 mmol/l arginine, pH 6.2, 30 mmol/l sodium sulfate, 3 % mannitol (w/v), 10 mM methionine, 0.01% pluronic F68 (w/v) and optionally 1 mmol/l CaCl_2 .

A preferred embodiment of the present invention refers to compositions comprising 10 to 10000 $\mu\text{g/ml}$ erythropoietin, preferably 25 to 2500 $\mu\text{g/ml}$ erythropoietin, and

- 15 a) 10 mM sodium/potassium phosphate, 100 mM NaCl, pH 7.0 or
b) 10 mM sodium phosphate, 120 mM sodium sulfate, pH 6.2 or
c) 10 mM sodium phosphate, 40 mM sodium sulfate, 3% mannitol (w/v) , pH 6.2 or
d) 10 mM sodium phosphate, 40 mM sodium sulfate, 3% mannitol (w/v), 10
20 mM methionine, 0.01% pluronic F68 (w/v), pH 6.2 or
e) 40 mM arginine, 30 mM sodium sulfate, 3% mannitol (w/v), pH 6.2 or
f) 40 mM arginine, 30 mM sodium sulfate, 3% mannitol (w/v), 10 mM methionine, 0.01% pluronic F68 (w/v), pH 6.2.

In the most preferred embodiment, the compositions comprise an amount
25 erythropoietin protein of 50, 100, 400, 800 or 2500 $\mu\text{g/ml}$. The most preferred compositions comprise either 10 mM sodium phosphate, 40 mM sodium sulfate, 3% mannitol (w/v), 10 mM methionine, 0.01% pluronic F68 (w/v), pH 6.2 or 40 mM arginine, 30 mM sodium sulfate, 3% mannitol (w/v), 10 mM methionine, 0.01% pluronic F68 (w/v), pH 6.2. Further details of such compositions are known from WO 01/87329.

30 The invention also relates to a method for treating disturbances of iron distribution in diabetes comprising administration of an effective amount of erythropoietin protein as defined above. Furthermore, the invention relates to a medicament for treating disturbances of iron distribution in diabetes characterized in that it contains an effective amount of erythropoietin protein. In the above cases, non insulin dependant diabetes
35 mellitus is the preferred form of diabetes.

The specific activity of EPO or EPO conjugates in accordance with this invention can be determined by various assays known in the art. The biological activity of the purified EPO proteins of this invention are such that administration of the EPO protein by injection to human patients results in bone marrow cells increasing production of reticulocytes and red blood cells compared to non-injected or control groups of subjects. The biological activity of the EPO proteins, or fragments thereof, obtained and purified in accordance with this invention can be tested by methods according to Annable, et al., Bull. Wld. Hlth. Org. (1972) 47: 99-112 and Pharm. Europa Spec. Issue Erythropoietin BRP Bio 1997(2). Another biological assay for determining the activity of EPO protein, the normocythaemic mouse assay, is described in the art (e.g. Pharm. Europa Spec. Issue Erythropoietin BRP Bio 1997(2), and the monography of erythropoietin of Ph. Eur. BRP.).

The invention will be better understood by reference to the following examples which illustrate but do not limit the invention described herein.

EXAMPLE

5


A middle-aged woman suffering from diabetes is checked for disturbances of iron distribution by determination of the following parameters - CRP (C reactive protein), ferritin and soluble transferrin receptor - as described P. Lehmann, M. Volkmann, J. Lotz, A. Baldauf, R. Roeddiger, poster presented at the AACC/CSCC, Annual Meeting, July 29 – August 2, 2001, Chicago, Illinois. The results show disturbances of iron distribution. The patient is treated subcutaneously with 150/U kg Recormon™ (commercially available erythropoietin protein) twice a week for a maximum of 12 weeks. Afterwards, determination of the parameters as described above shows an improvement of the disorder of iron deficiency.

CLAIMS

1. The use of erythropoietin protein in the manufacture of a medicament for the treatment of disturbances of iron distribution in diabetes.
- 5 2. The use of claim 1, wherein diabetes is non insulin dependant diabetes mellitus.
3. The use according to any of claims 1 to 2, wherein the erythropoietin protein is a human erythropoietin.
4. The use according to claim 3, wherein the erythropoietin protein is epoetin alfa or epoetin beta.
- 10 5. The use according to any of claims 1 to 4, wherein the erythropoietin protein is expressed by endogenous gene activation.
6. The use according to any of claims 1 to 2, wherein the erythropoietin protein has the amino acid sequence of SEQ ID NO:1 or SEQ ID NO:2.
- 15 7. The use according to any of claims 1 to 2, wherein the erythropoietin protein has the sequence of human erythropoietin modified by the addition of from 1 to 6 glycosylation sites.
8. The use of according to any of claims 1 to 2, wherein the erythropoietin protein is darbepoetin alfa.
- 20 9. The use according to any of claims 1 to 2, wherein the erythropoietin protein as defined in any of claims 3 to 8 is pegylated.
- 25 10. The use according to claim 9, wherein the erythropoietin protein is a conjugate, said conjugate comprising an erythropoietin protein having at least one free amino group and having the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells and selected from the group consisting of human erythropoietin and analogs thereof which have a sequence of human erythropoietin modified by the addition of from 1 to 6 glycosylation sites or a rearrangement of at least one glycosylation site; said erythropoietin protein being covalently linked to n poly(ethylene glycol) groups of the formula $-\text{CO}-(\text{CH}_2)_x-(\text{OCH}_2\text{CH}_2)_m-\text{OR}$ with the $-\text{CO}$ of each
30 poly(ethylene glycol) group forming an amide bond with one of said amino


groups; wherein R is lower-alkyl; x is 2 or 3; m is from about 450 to about 900; n is from 1 to 3; and n and m are chosen so that the molecular weight of the conjugate minus the erythropoietin protein is from 20 kilodaltons to 100 kilodaltons.

- 5 11. The use according to claim 10, wherein x is 2, m is 650 to 750, n is 1 and R is methyl.
- 10 12. The use according to claim 9, wherein the erythropoietin protein is a conjugate, said conjugate comprising an erythropoietin protein having at least one free amino group and having the *in vivo* biological activity of causing bone marrow cells to increase production of reticulocytes and red blood cells and selected from the group consisting of human erythropoietin protein and analogs thereof which have the primary structure of human erythropoietin protein modified by the addition of from 1 to 6 glycosylation sites; said erythropoietin protein being covalently linked to from one to three lower-alkoxy poly(ethylene glycol)
- 15 groups, each poly(ethylene glycol) group being covalently linked to the erythropoietin protein *via* a linker of the formula $-C(O)-X-S-Y-$ with the C(O) of the linker forming an amide bond with one of said amino groups, X is $-(CH_2)_k-$ or $-CH_2(O-CH_2-CH_2)_k-$, k is from 1 to 10, Y is



- 20 the average molecular weight of each poly(ethylene glycol) moiety is from about 20 kilodaltons to about 40 kilodaltons, and the molecular weight of the conjugate is from about 51 kilodaltons to about 175 kilodaltons.

- 25 13. The use according to claim 12, with an erythropoietin conjugate of the formula:



wherein n is an integer from 1 to 3; m is an integer from 450 to 900; R is lower-alkyl; X is $-(\text{CH}_2)_k-$ or $-\text{CH}_2(\text{O}-\text{CH}_2-\text{CH}_2)_k-$, k is 1 to 10 and P is the residue of the erythropoietin protein without the n amino groups which form an amide linkage with X.

5

14. A method for treating disturbances of iron distribution in diabetes comprising administration of an effective amount of erythropoietin protein.

15. A medicament for treating disturbances of iron distribution in diabetes characterized in that it contains an effective amount of erythropoietin protein.

10

16. The invention as hereinbefore defined.

67
29 Aug. 2002

ABSTRACT

- 26 -

The present invention relates to the use of erythropoietin for the treatment of disturbances of iron distribution in diabetes.

5

FIGURES

Figure 1

Ala Pro Pro Arg Leu Ile Cys Asp Ser Arg¹⁰ Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys²⁰
 Glu Ala Glu Asn Ile Thr Thr Gly Cys Ala³⁰ Glu His Cys Ser Leu Asn Glu Asn Ile Thr⁴⁰
 Val Pro Asp Thr Lys Val Asn Phe Tyr Ala⁵⁰ Trp Lys Arg Met Glu Val Gly Gln Gln Ala⁶⁰
 Val Glu Val Trp Gln Gly Leu Ala Leu Leu⁷⁰ Ser Glu Ala Val Leu Arg Gly Gln Ala Leu⁸⁰
 Leu Val Asn Ser Ser Gln Pro Trp Glu Pro⁹⁰ Leu Gln Leu His Val Asp Lys Ala Val Ser¹⁰⁰
 Gly Leu Arg Ser Leu Thr Thr Leu Leu Arg¹¹⁰ Ala Leu Gly Ala Gln Lys Glu Ala Ile Ser¹²⁰
 Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu¹³⁰ Arg Thr Ile Thr Ala Asp Thr Phe Arg Lys¹⁴⁰
 Leu Phe Arg Val Tyr Ser Asn Phe Leu Arg¹⁵⁰ Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala¹⁶⁰
 Cys Arg Thr Gly Asp¹⁶⁵

Figure 2

Ala Pro Pro Arg Leu Ile Cys Asp Ser Arg¹⁰ Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys²⁰
 Glu Ala Glu Asn Ile Thr Thr Gly Cys Ala³⁰ Glu His Cys Ser Leu Asn Glu Asn Ile Thr⁴⁰
 Val Pro Asp Thr Lys Val Asn Phe Tyr Ala⁵⁰ Trp Lys Arg Met Glu Val Gly Gln Gln Ala⁶⁰
 Val Glu Val Trp Gln Gly Leu Ala Leu Leu⁷⁰ Ser Glu Ala Val Leu Arg Gly Gln Ala Leu⁸⁰
 Leu Val Asn Ser Ser Gln Pro Trp Glu Pro⁹⁰ Leu Gln Leu His Val Asp Lys Ala Val Ser¹⁰⁰
 Gly Leu Arg Ser Leu Thr Thr Leu Leu Arg¹¹⁰ Ala Leu Gly Ala Gln Lys Glu Ala Ile Ser¹²⁰
 Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu¹³⁰ Arg Thr Ile Thr Ala Asp Thr Phe Arg Lys¹⁴⁰
 Leu Phe Arg Val Tyr Ser Asn Phe Leu Arg¹⁵⁰ Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala¹⁶⁰
 Cys Arg Thr Gly Asp Arg¹⁶⁶

23. Aug. 2002

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT:

10

- (A) NAME: F.Hoffmann-La Roche AG
- (B) STREET: 124 Grenzacherstrasse
- (C) CITY: Basle
- (E) COUNTRY: Switzerland
- (F) POSTAL CODE (ZIP): CH-4070
- (G) TELEPHONE: (61) 688 11 11
- (H) TELEFAX: (61) 688 13 95
- (I) TELEX: 962 292 hlr ch

15

(ii) TITLE OF INVENTION: New Pharmaceutical Composition

(iii) NUMBER OF SEQUENCES: 2

20

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: WORD
- (D) SOFTWARE: PatentIn Release 2.0

25

<170> PatentIn Ver. 2.0

30

<210> 1
<211> 165
<212> PRT
<213> Homo sapiens

35

<400> 1
Ala Pro Pro Arg Leu Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu
1 5 10 15

40

Leu Glu Ala Lys Glu Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His
20 25 30

Cys Ser Leu Asn Glu Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe
35 40 45

45

Tyr Ala Trp Lys Arg Met Glu Val Gly Gln Gln Ala Val Glu Val Trp
50 55 60

50

Gln Gly Leu Ala Leu Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu
65 70 75 80

Leu Val Asn Ser Ser Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp
85 90 95

55

Lys Ala Val Ser Gly Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu
100 105 110

Gly Ala Gln Lys Glu Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala
115 120 125

60

Pro Leu Arg Thr Ile Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val

	130		135		140	
	Tyr Ser Asn Phe Leu Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala					
	145		150		155	160
5	Cys Arg Thr Gly Asp					
			165			
10	<210> 2					
	<211> 166					
	<212> PRT					
	<213> Homo sapiens					
15	<400> 2					
	Ala Pro Pro Arg Leu Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu					
	1		5		10	15
20	Leu Glu Ala Lys Glu Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His					
			20		25	30
	Cys Ser Leu Asn Glu Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe					
			35		40	45
25	Tyr Ala Trp Lys Arg Met Glu Val Gly Gln Gln Ala Val Glu Val Trp					
			50		55	60
	Gln Gly Leu Ala Leu Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu					
			65		70	75
30	Leu Val Asn Ser Ser Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp					
			85		90	95
35	Lys Ala Val Ser Gly Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu					
			100		105	110
	Gly Ala Gln Lys Glu Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala					
			115		120	125
40	Pro Leu Arg Thr Ile Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val					
			130		135	140
	Tyr Ser Asn Phe Leu Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala					
			145		150	155
45	Cys Arg Thr Gly Asp Arg					
			165			

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.